arXiv:0901.3015v1 [math.AC] 20 Jan 2009

RESOLUTION OF UNMIXED BIPARTITE GRAPHS

FATEMEH MOHAMMADI AND SOMAYEH MORADI

Abstract. For an unmixed bipartite graph G we consider the lattice of vertex covers $\mathcal{L}_G$ and compute depth, projective dimension and extremal Betti-numbers of $R/I(G)$ in terms of this lattice.

Introduction

In recent years edge ideals of graphs have been intensively studied. One central question in this context is when the edge ideal is Cohen-Macaulay. In the case of chordal graphs and bipartite graphs a nice and complete answer is known, see [6] and [7]. In particular, in these cases one knows the projective dimension of the edge ideal. For bipartite Cohen-Macaulay graphs the Cohen-Macaulay type is also known. There are also several papers in which the regularity of edge ideals has been studied, (see [2], [3], [11]). It is impossible to classify all Cohen-Macaulay graphs or to give uniform formulas for the projective dimension or the regularity for all graphs. Nevertheless there are interesting classes of graphs where the homological data of the edge ideals can be described. In this paper we consider unmixed bipartite graphs and study the resolution of their edge ideals.

To be more specific, let G be a finite graph with vertex set $V(G) = \{v_1, \dots, v_n\}$ and edge set $E(G)$. Throughout we assume that G has no isolated vertices. The graph G is called bipartite, if $V(G) = X \cup Y$ with $X \cap Y = \emptyset$ such that $E(G) \subseteq X \times Y$.

A vertex cover of G is a subset C of $V(G)$ such that each edge has at least one vertex in C. A minimal vertex cover C of G is a vertex cover such that no proper subset of C is a vertex cover of G. The graph G is called unmixed if all its minimal vertex covers are of the same cardinality.

Let G be an unmixed bipartite graph with vertex set $V(G) = \{x_1, \dots, x_n\} \cup \{y_1, \dots, y_m\}$. Since G is unmixed, it follows that $n = m$. For any unmixed bipartite graph there is a perfect matching, see [10]. Therefore we may assume that $\{x_i, y_i\}$ is an edge of G for all i. So each minimal vertex cover of G is of the form $\{x_{i_1}, \dots, x_{i_s}, y_{i_{s+1}}, \dots, y_{i_n}\}$, where $\{i_1, \dots, i_n\} = [n]$. For any minimal vertex cover C of G let $\bar{C} = C \cap \{x_1, \dots, x_n\}$. Let B_n be the Boolean lattice on the set $\{x_1, \dots, x_n\}$. In [5] it is shown that for any unmixed bipartite graph G, the subset

$$\mathcal{L}_G = \{\bar{C} : C \text{ is a minimal vertex cover of } G\} \subseteq \{x_1, \dots, x_n\}$$

is a sublattice of B_n which contains $\emptyset$ and X, and for any such sublattice $\mathcal{L}$ of B_n, there exists an unmixed bipartite graph G such that $\mathcal{L} = \mathcal{L}_G$. In our description of the resolution of the edge ideal of an unmixed bipartite graph, we use in a substantial way this characterization of such graphs in terms of their vertex cover lattices.

1991 *Mathematics Subject Classification.* 13D02, 13P10, 13D40, 13A02.

In the following let $\mathcal{L}_G$ be a sublattice of the Boolean lattice B_n corresponding to the unmixed bipartite graph G. Therefore $\wedge$ and $\vee$ in $\mathcal{L}_G$ are just taking the intersection and union. Attached to the lattice $\mathcal{L}_G$ is a monomial ideal $H_{\mathcal{L}_G}$ in the polynomial ring $K[x_1, \dots, x_n, y_1, \dots, y_n]$, defined as follows: for each element $p \in \mathcal{L}_G$, let $u_p = X_p Y_{[n]\setminus p}$, where $X_p = \prod_{i \in p} x_i$ and $Y_{[n]\setminus p} = \prod_{j \in [n]\setminus p} y_j$. Then $H_{\mathcal{L}_G}$ is generated by the monomials u_p, $p \in \mathcal{L}_G$. The edge ideal $I(G)$ which we are interested in is just the Alexander dual of $H_{\mathcal{L}_G}$. Thus we may apply the Bayer-Charalambous-Popescu theory [1, Theorem 2.8] which relates the multigraded extremal Betti-numbers of $H_{\mathcal{L}_G}$ and $I(G)$.

In Section 1 we describe the multigraded minimal free resolution of the monomial ideal $H_{\mathcal{L}_G}$. In fact, the resolution we describe in Theorem 1.2 is a variation of the resolution of Hibi ideals of meet-distributive semilattices, given in [7, Theorem 2.1]. In Theorem 1.3 we also describe the differentials of this resolution. This information is not needed later. More important is the fact, that the multigraded basis elements of the resolution, can be identified with the Boolean sublattices of $\mathcal{L}_G$, see Proposition 1.4. Having this identification, it turns out that the multigraded extremal Betti-numbers correspond to the maximal Boolean sublattices of $\mathcal{L}_G$. In Section 2 we use Alexander duality and Theorem in [1] to obtain the multigraded extremal Betti-numbers of $I(G)$. With this information at hand, we can express the depth and regularity of $R/I(G)$ in terms of the lattice $\mathcal{L}_G$, see Corollary 1.6. As a further corollary we obtain a lower bound for the last nonzero total Betti-number of $R/I(G)$. We do not know of any example where this lower is not achieved. It would be always achieved if one could prove the following: all nonzero multigraded Betti-number in the last step of the resolution are extremal (in the multigraded sense). There is a simple argument, given in the proof of Proposition 1.10 that whenever I is monomial Cohen-Macaulay ideal, then all multigraded extremal Betti-numbers appear at the end of the resolution of I.

After having finished the paper, Professor Herzog informed us that Kummini in his thesis and in a preprint paper has also computed the depth and regularity of unmixed bipartite graphs, see [9]. His approach and the terms in which he expresses these invariants differ from ours.

1. Minimal Free Resolution of $\mathcal{L}_G$

The purpose of this section it to construct a resolution of the ideal $H_{\mathcal{L}_G}$ which is a modification of the resolution given [7, Theorem 2.1] adopted to our situation. The differences between our resolution and the one given in [7] arises from the fact that the lattices under consideration are differently embedded into Boolean lattice. This fact is important, because the multidegrees of the resolution depends on the embedding.

In order to guarantee the minimality of the resolution which we are going to describe we need the following result.

Lemma 1.1. *Let $p \in \mathcal{L}_G$. For any two distinct subsets $S, S' \subseteq N(p)$, we have $\wedge\{q; q \in S\} \neq \wedge\{q; q \in S'\}$.*

Proof. Let $S, S' \subseteq N(p)$ be distinct subsets. One can assume that $S \nsubseteq S'$. Let $q_1 \in S \setminus S'$. If $\wedge\{q; q \in S\} = \wedge\{q; q \in S'\}$, then $(\wedge\{q; q \in S\}) \vee q_1 = (\wedge\{q; q \in S'\}) \vee q_1$. Since $\mathcal{L}_G$ is distributive $\wedge(\{q \vee q_1; q \in S\}) = \wedge(\{q \vee q_1; q \in S'\})$. For any $q \in S'$,

$q \vee q_1 = p$, therefore $\wedge(\{q \vee q_1; q \in S'\}) = p$. But then $\wedge(\{q \vee q_1; q \in S\}) = q_1$, a contradiction. $\square$

From the above lemma it is easy to see that for any subsets S and S' of $N(p)$ with $S \subseteq S'$, we have $|S'| - |S| \leq |\wedge\{q; q \in S\}| - |\wedge\{q; q \in S'\}|$. In the following we denote by $\hat{0}$ and $\hat{1}$ the minimal and maximal element of $\mathcal{L}_G$. For any $p \in \mathcal{L}$, rank of p which is denoted by $\operatorname{rank}(p)$, is the maximal length of chains descending from p. We extend the partial rank order on $\mathcal{L}_G$ to a total order $\prec$.

Theorem 1.2. *There exists a minimal multigraded free resolution $\mathbb{F}$ of $H_{\mathcal{L}_G}$ such that for each $i \geq 0$, the free module $\mathbb{F}_i$ has a basis with basis elements $b(p; S)$, where $p \in \mathcal{L}_G$ and S is a subset of the set of lower neighbors $N(p)$ of p with $|S| = i$ and multidegree of $b(p; S)$ is the least common multiple of u_p and all u_q with $q \in S$.*

Proof. The construction of resolution is as in the proof of [7, Theorem 2.1] by mapping cone. For any $p \in \mathcal{L}_G$ we construct inductively a complex $\mathbb{F}(p)$ which is a multigraded free resolution of the ideal $H_{\mathcal{L}_G}(p) = \langle u_q; q \preceq p\rangle$. The complex $\mathbb{F}(\hat{0})$ is defined as $\mathbb{F}_i(\hat{0}) = 0$ for $i > 0$ and $\mathbb{F}_0(\hat{0}) = S$. Now, let $p \in \mathcal{L}_G$ and $q \in \mathcal{L}_G$, $q \prec p$ be the element preceding p. Then $H_{\mathcal{L}_G}(p) = (H_{\mathcal{L}_G}(q), u_p)$, and hence we have the exact sequence of multigraded S-modules

$$0 \longrightarrow (S/L)(-\operatorname{multideg} u_p) \longrightarrow S/H_{\mathcal{L}_G}(q) \longrightarrow S/H_{\mathcal{L}_G}(p) \longrightarrow 0,$$

where L is the colon ideal $H_{\mathcal{L}_G}(q) : u_p$. Let $u_{q'}/[u_{q'}, u_p] \in L$, where $q' < q$ and let $t \in [q' \wedge q, p] \cap N(p)$. Then $u_t/[u_t, u_p]$ divides $u_{q'\wedge p}/[u_{q'\wedge p}, u_p]$ and $u_{q'\wedge p}/[u_{q'\wedge p}, u_p]$ divides $u'_q/[u'_q, u_p]$. Therefore we have $L = (\{u_t/[u_t, u_p]\}_{t \in N(p)})$.

Let $\mathbb{T}$ be the Taylor complex associated with the sequence $u_t/[u_t, u_p]$, $t \in N(p)$, where the order of the sequence is given by the order $\prec$ on the elements of $\mathbb{L}$. Then T_i has a basis with elements $e_{t_1} \wedge e_{t_2} \wedge \cdots \wedge e_{t_i}$, where $t_1 \prec t_2 \prec \cdots \prec t_i$. The multidegree of $e_{t_1} \wedge e_{t_2} \wedge \cdots \wedge e_{t_i}$ is the least common multiple of the elements $u_{t_j}/[u_{t_j}, u_p]$ for $j = 1, \ldots, i$. The shifted complex $\mathbb{T}(-\operatorname{multideg}(u_p))$ is a multigraded free resolution of $(S/L)(-\operatorname{multideg}(u_p))$. Let $b(p; t_1, \ldots, t_i)$ be basis element of $\mathbb{T}_i(-\operatorname{multideg}(u_p))$ corresponding to $e_{t_1} \wedge e_{t_2} \wedge \cdots \wedge e_{t_i}$. Then $\operatorname{multideg}(b(p; t_1, \ldots, t_i))$=$\operatorname{multideg}(u_p)$+$\operatorname{multideg}(e_{t_1} \wedge e_{t_2} \wedge \cdots \wedge e_{t_i}) = lcm(u_p, u_{t_1}, \ldots, u_{t_i})$. This resolution is minimal since for any $t_1 \prec t_2 \prec \cdots \prec t_i$ we have $lcm(f_1, \ldots, f_i)/lcm(f_1, \ldots, \check{f}_j, \ldots, f_i) = Y_{p \setminus \bigwedge_{l=1}^{i} t_l} / Y_{p \setminus \bigwedge_{l=1, l \neq j}^{i} t_l} \neq 1$.

The monomorphism $(S/L)(-\operatorname{multideg}(u_p)) \longrightarrow S/H_{\mathcal{L}_G}(q)$ induces a comparison map $\alpha : \mathbb{T}(-\operatorname{multideg}(u_p)) \longrightarrow \mathbb{F}(q)$ of multigraded complexes. Let $\mathbb{F}(p)$ be the mapping cone of α. Then $\mathbb{F}(p)$ is a multigraded free S-resolution of $H_{\mathcal{L}_G}(p)$ with the desired multigraded basis.

We claim that this resolution is minimal. For any two basis elements $b(p; S)$ and $b(q; T)$ with $|T| = |S| - 1$. We show that the coefficient of $b(q; T)$ in $\partial b(p; S)$ is either zero or a monomial $\neq 1$. First assume that $p = q$. If $T \subseteq S$, then the coefficient is $\operatorname{multideg}(b(p; S))/\operatorname{multideg}(b(p, T)) = Y_A$, where $A = \wedge\{r; r \in T\} \setminus \wedge\{r; r \in S\}$. Since A is a nonempty set by Lemma 1.1, then $Y_A \neq 1$. If $T \nsubseteq S$ and $\operatorname{multideg}(b(p; T))$ divides $\operatorname{multideg}(b(p; S))$, then $\wedge\{r; r \in S\} \leq \wedge\{r; r \in T\}$. Therefore $\wedge\{r; r \in N(p)\} = \wedge\{r; r \in N(p) \setminus (T \setminus S)\}$, which is a contradiction by Lemma1.1. Now, assume that $q < p$. If $\operatorname{multideg}(b(q; T))$ divides $\operatorname{multideg}(b(p; S))$, then the coefficient is $X_{p \setminus q} Y_B$ for some set $B \subseteq [n]$ and so it is not 1. In the remaining case $q \nless p$, $\operatorname{multideg}(b(q; T))$ does not divide $\operatorname{multideg}(b(p; S))$. $\square$

In the following theorem we describe the maps in the resolution of Theorem 1.2. These information is just for curiosity and is not needed to get the results which come after. Before describing the maps in the resolution we fix some notation. For a subset $S \subset \mathcal{L}_G$ and $q \in S$ let $\sigma(q;S) = |\{r \in S; r \prec q\}|$. Let $X_{p\setminus q} = \prod_{i\in p\setminus q} x_i$ and $Y_{S_q} = \prod_{i\in(\bigwedge_{r\in S\setminus\{q\}} r)\setminus q} y_i$.

Theorem 1.3. *For any $p \in \mathcal{L}_G$ and $S \subseteq N(p)$ we have*

$$\partial(b(p;S)) = \sum_{q\in S}(-1)^{\sigma(q;S)}(Y_{S_q} b(p; S\setminus\{q\}) - X_{p\setminus q} b(q; q\wedge(S\setminus\{q\}))).$$

Proof. First we remind that for any q_i and q_j in $N(p)$ we have $q_i \wedge q_j \in N(q_i)$, since $[q_i \wedge q_j, p]$ is isomorphic to a Boolean lattice of rank two. Then we have $q\wedge(S\setminus\{q\}) \subseteq N(q)$. We have $\operatorname{multideg}(b(p;S)) = X_p Y_{[n]\setminus(\bigwedge_{r\in S} r)}$ and $\operatorname{multideg}(b(p;S\setminus\{q\})) = X_p Y_{[n]\setminus(\bigwedge_{r\in S\setminus q} r)}$. Also, $\{i; i \notin \bigwedge_{r\in S} r\} \setminus \{i; i \notin \bigwedge_{r\in S\setminus q} r\} = \{i; i \in \bigwedge_{r\in S\setminus q} r \setminus q\}$ shows that $Y_{[n]\setminus(\bigwedge_{r\in S} r)}/Y_{[n]\setminus(\bigwedge_{r\in S\setminus q} r)} = Y_{S_q}$ and so $\operatorname{multideg}(b(q; q\wedge(S\setminus\{q\}))) = X_q Y_{[n]\setminus\bigwedge_{r\in S\setminus q}(q\wedge r)} = X_q Y_{[n]\setminus\bigwedge_{r\in S}(r)}$ $=\operatorname{multideg}(b(p;S))/X_{p\setminus q}$. Therefore, ∂ is a multi-homogeneous differential. Consider the mapping cone constructed in Theorem 1.2. The differential given by that mapping cone is

$$\partial_i = (\partial_i^{\mathbb{T}} + (-1)^i\alpha_i, \partial_{i+1}^{\mathbb{F}(q)}) \text{ for all } i.$$

Comparing this equation with ∂ which is defined in theorem it is enough to show that for any $S \subseteq N(p)$ we have:

(i) $\partial^{\mathbb{T}}(b(p;S)) = \sum_{q\in S}(-1)^{\sigma(q,S)} Y_{S_q} b(p; S\setminus\{q\})$, which is exactly the definition of Taylor complex.

(ii) We can choose α such that

$$(-1)^i\alpha_i(b(p;S)) = -\sum_{q\in S}(-1)^{\sigma(q,S)} X_{p\setminus q} b(q; q\wedge S\setminus\{q\}).$$

We show that $\alpha : \mathbb{C} \to \mathbb{F}(q)$ is a complex homomorphism. So it is enough to show that for any $b(p;S) \in T_i$

$$\partial_i^{F(q)}\alpha_i(b(p;S)) = \alpha_{i-1}\partial_i^{\mathbb{T}}(b(p;S)).$$

First we compute the left side of the above equation. Then we have

(1) $\partial_i^{F(q)}\alpha_i(b(p;s)) = (-1)^{i+1}\sum_{q\in S}(-1)^{\sigma(q;S)} X_{p\setminus q}\partial_i^{F(q)}(b(q; q\wedge(S\setminus\{q\})))$.

By induction hypothesis we have:

$$\begin{aligned}(2)\quad \partial_i^{F(q)}(b(q; q\wedge(S\setminus\{q\}))) \quad &= \quad \sum_{q'\in S\setminus q}(-1)^{\sigma(q';S\setminus\{q\})} Y_{(q\wedge S)_{q\wedge q'}} b(q; q\wedge(S\setminus\{q,q'\})) \\ &- \quad X_{q\setminus(q\wedge q')} b(q\wedge q'; q'\wedge[(q\wedge(S\setminus\{q\})\setminus\{q,q'\}])).\end{aligned}$$

Since $q \vee q' = p$ for any $q, q' \in N(p)$ we have $q\setminus(q\wedge q') = p\setminus q'$. Also from $\bigwedge_{r\in S\setminus\{q,q'\}}\setminus\{q,q'\} = \bigwedge_{r\in S\setminus\{q'\}}\setminus\{q,q'\} = \bigwedge_{r\in S\setminus\{q'\}}\setminus\{q'\}$ we see that $Y_{(q\wedge S)_{q\wedge q'}} = Y_{S_{q'}}$. Considering these statements and substituting (2) in (1) we get

(3) $\partial_i^{F(q)}\alpha_i(b(p;s)) =$

$(-1)^{i+1}\sum_{q,q'\in S,q\neq q'}(-1)^{\sigma(q;S)+\sigma(q';S\setminus\{q\})}X_{p\setminus q}Y_{S_{q'}}b(q;q\wedge(S\setminus\{q,q'\}))$.

On the other hand we have

$$(4)\quad \alpha_{i-1}\partial_i^{\mathbb{T}}(b(p;S))=(-1)^{\sigma(q;S)}Y_{S_q}\alpha_{i-1}(b(p;S\setminus\{q\}))$$

After substituting $\alpha_{i-1}(b(p;S\setminus\{q\}))$ in (4) and using $q\setminus(q\wedge q')=p\setminus q'$ we get

$$(5)\quad \alpha_{i-1}\partial_i^{\mathbb{T}}(b(p;S))=$$

$(-1)^{i+1}\sum_{q,q'\in S,q\neq q'}(-1)^{\sigma(q;S)+\sigma(q';S\setminus\{q\})}Y_{S_q}X_{p\setminus q'}b(q';q'\wedge(S\setminus\{q,q'\}))$.

After exchanging q and q' we see that the equations (3) and (5) are equal, which completes the proof. □

The next observation is of crucial importance for understanding the i-extremal and extremal Betti-numbers of $H_{\mathcal{L}_G}$ and $I(G)$.

Proposition 1.4. *There is a correspondence between the basis elements $b(p;S)$ and intervals in $\mathcal{L}_G$, which are isomorphic to a Boolean lattice.*

Proof. For any basis element $b(p;S)$, we show that the interval $[\wedge\{q;q\in S\},p]$ is isomorphic to $B_{|S|}$. Let $S=\{q_1,\dots,q_n\}$ and $v_i=\wedge\{q;q\in S\setminus\{q_i\}\}$. The interval $[\wedge\{q;q\in S\},p]$ is a Boolean lattice on $v_1,\dots,v_n$.

For any $I\in[\wedge\{q;q\in S\},p]$, let $i_1,\dots,i_k$ be the indices that $I\leq q_{i_j}$, $1\leq j\leq k$. Then $I\leq\wedge\{q_{i_j};1\leq j\leq k\}$. If $I\neq\wedge\{q_{i_j};1\leq j\leq k\}$, then there exists $x\in\wedge\{q_{i_j};1\leq j\leq k\}\setminus I$. Since $\wedge\{q;q\in S\}\leq I$, there exists $1\leq l\leq n$, $l\neq i_1,\dots,i_k$ such that $x\notin q_l$. But then $x\notin I\vee q_l=p$, a contradiction. Thus $I=\wedge\{q_{i_j};1\leq j\leq k\}=\vee\{v_j;1\leq j\leq n,j\neq i_1,\dots,i_k\}$. Let Φ be a function from the set of basis elements to intervals in $\mathcal{L}_G$, which are isomorphic to a Boolean lattice such that $\Phi(b(p;S))=[\wedge\{q;q\in S\},p]$. From Lemma 1.1 we know that Φ is a monomorphism. For any interval $[J,I]$ in $\mathcal{L}_G$ isomorphic to a Boolean lattice, set $S=N(I)\wedge[J,I]$. Then $[J,I]=[\wedge\{q;q\in S\},I]$ and Φ is surjective. □

In the following we denote by A_G the set of elements $p\in\mathcal{L}_G$ such that the interval $[\wedge\{r;r\in N(p)\},p]$ is isomorphic to a maximal Boolean lattice in $\mathcal{L}_G$.

Lemma 1.5. *Let $p\in A_G$. For any $q\in\mathcal{L}_G$ such that $[\wedge\{r;r\in N(q)\},q]\subseteq[\wedge\{r;r\in N(p)\},p]$, we have*

$$|q|-|N(q)|-|\wedge\{r;r\in N(q)\}|\leq|p|-|N(p)|-|\wedge\{r;r\in N(p)\}|.$$

Proof. Let $[\wedge\{r;r\in N(p)\},p]$ be a Boolean lattice on the elements $v_1,\dots,v_n$. Then $|N(p)|=n$, $|N(q)|\leq n$ and $v_i\wedge v_j=\wedge\{r;r\in N(p)\}$ for any $1\leq i<j\leq n$. Also $v_1\vee\cdots\vee v_n=p$. Without loss of generality assume that $q=v_1\vee\cdots\vee v_m$ for some $m<n$. Then $|N(q)|=m$. We claim that for any $1\leq i\leq n$, there exists an element $x_i\in v_i$ such that x_i is not in any other v_j, $1\leq j\leq n$. Otherwise let $1\leq i\leq n$ be such that for any $x\in v_i$, there exists $j\neq i$ with $x\in v_j$. It means that $v_i\leq\wedge\{r;r\in N(p)\}$, a contradiction. Thus $|p|-|q|\geq n-m$. Since $|\wedge\{r;r\in N(p)\}|\leq|\wedge\{r;r\in N(q)\}|$, we get the inequality. □

As a first corollary we obtain

Corollary 1.6. *Let G be an unmixed bipartite graph on (X, Y) such that $|X| = |Y| = n$ and $A_G \subseteq \mathcal{L}_G$ be the set defined above. Then*

(i) $\operatorname{depth}(R/I(G)) = n - \max_{p \in A_G}\{|p| - |N(p)| - |\wedge\{r; r \in N(p)\}|\}$;

(ii) $\operatorname{reg}(R/I(G)) = \max_{p \in \mathcal{L}_G} |N(p)|$.

Proof. For any basis element $b(p; S)$ in $\mathbb{F}_{|S|}$, one has $\operatorname{multideg}(b(p; S)) = X_p Y_{[n] \setminus \wedge\{r; r \in S\}}$ and so $\deg(b(p; S)) = |p| + n - |\wedge\{r; r \in S\}|$. For $S \subseteq N(p)$ from Lemma 1.1 we have $|\wedge\{r; r \in N(p)\}| + |N(p)| \leq |\wedge\{r; r \in S)\}| + |S|$. Therefore by Lemma 1.5

$$\operatorname{reg}\ (H_{\mathcal{L}_G}) = n + \max_{p \in A_G}\{|p| - |N(p)| - |\wedge\{r; r \in N(p)\}|\}.$$

Since $\operatorname{reg}(H_{\mathcal{L}_G}) = \operatorname{pd}(R/I(G))$, from Auslander-Buchsbaum formula one has (i). The other statement is clear from Theorem 1.2. □

Recall that a multigraded Betti-number $\beta_{i,b}$ is called i-extremal if $\beta_{i,c} = 0$ for all $c > b$, that is all multigraded entries below b on the i-th column vanish in the Betti diagram as a Macaulay output [Mac]. Define $\beta_{i,b}$ to be extremal if $\beta_{j,c} = 0$ for all $j \geq i$, and $c > b$ so $|c| - |b| \geq j - i$. In other words, $\beta_{i,b}$ corresponds to the top left corner of a box of zeroes in the multigraded Betti diagram. A graded Betti-number $\beta_{i,r}$ is called extremal if $\beta_{j,l} = 0$ for any $j \geq i$, $l > r$ and $l - r \geq j - i$. In other words, $\beta_{i,r}$ corresponds to the top left corner of a box of zeroes in the multigraded Betti diagram.

Corollary 1.7. *All multigraded Betti-numbers in homological degree i are i-extremal. A multigraded Betti-number $\beta_{i,b}$ of $H_{\mathcal{L}_G}$ is extremal if and only if there exists $p \in A_G$ such that $i = |N(p)|$ and $b = \operatorname{multideg}(b(p; N(p)))$.*

Proof. Let $b(p; S)$ and $b(p'; S')$ be two basis elements of $\mathbb{F}_i$ such that $\operatorname{multideg}(b(p; S))$ divides $\operatorname{multideg}(b(p'; S'))$. Then $p \leq p'$ and $\wedge\{q; q \in S'\} \leq \wedge\{q; q \in S\}$ and $|S| = |S'| = i$. Therefore $[\wedge\{q; q \in S\}, p] \subseteq [\wedge\{q; q \in S'\}, p']$. Since both intervals are isomorphic to a Boolean lattice of rank i, one has $p = p'$ and $\wedge\{q; q \in S\} = \wedge\{q; q \in S'\}$. Since $S, S' \subseteq N(p)$ from Lemma 1.1 we have $S = S'$. Therefore all multigraded Betti-numbers in homological degree i are i-extremal.

For any basis element $b(p; S)$, $\operatorname{multideg}(b(p; S)) = X_p Y_{[n] \setminus \wedge\{q; q \in S\}}$. Therefore $\operatorname{multideg}(b(p; S))$ divides $\operatorname{multideg}(b(p; N(p)))$ and $|N(p)| - |S| \leq |\wedge\{q; q \in S\}| - |\wedge\{q; q \in N(p)\}|$. Therefore $\beta_{i,b}$ is extremal only if $b = \operatorname{multideg}(p; N(p))$ for some $p \in \mathcal{L}_G$. Let p and q be two elements of $\mathcal{L}_G$ and $\operatorname{multideg}(b(p; N(p)))$ divides $\operatorname{multideg}(b(q; N(q)))$. Then $p \leq q$ and $\wedge\{r; r \in N(q)\} \leq \wedge\{r; r \in N(p)\}$. Therefore $[\wedge\{r; r \in N(p)\}, p] \subseteq [\wedge\{r; r \in N(q)\}, q]$. Then $\beta_{|N(p)|,b}$ is extremal precisely when $b = \operatorname{multideg}(b(p; N(p)))$, where $[\wedge\{q; q \in N(p)\}, p]$ is a maximal Boolean sublattice of $\mathcal{L}_G$. □

The graded extremal Betti-numbers of $R/I(G)$ can be seen from the lattice $\mathcal{L}_G$. Indeed, the Betti-number $\beta_{i,i+j}$ for $R/I(G)$ is extremal if and only if there exists $p \in A_G$ with $|N(p)| = j$ and $n + |p| - |N(p)| - |\wedge\{r; r \in N(p)\}| = i$ such that:

(a) For any $q \in A_G$ with $|N(q)| > |N(p)|$, one has

$$|q| - |N(q)| - |\wedge\{r; r \in N(q)\}| < |p| - |N(p)| - |\wedge\{r; r \in N(p)\}|.$$

(b) For any $q \in A_G$ with $|N(q)| = |N(p)|$, one has

$$|q| - |\wedge\{r; r \in N(q)\}| \leq |p| - |\wedge\{r; r \in N(p)\}|.$$

This statement follows immediately from the definition of graded extremal Betti-numbers by using [1, Theorem 2.8].

Example 1.8. For any $p \in \mathcal{L}_G$ let $f(p) = |p| - |N(p)| - |\wedge\{r; r \in N(p)\}|$. In the following lattice we have $f(p) = 1$ for $p = \{1,2,3\}$, $p = \{1,2,3,4\}$ and $p = \{1,2,3,4,5,6,7\}$ and $f(p) = 0$ for $p = \{1,2,3,5\}$ and $p = \{1,2,3,4,5\}$. Therefore by the statement above, the extremal Betti-number of the unmixed bipartite graph corresponding to this lattice is $\beta_{i,i+j}$ for $i = 7 + f(\{1,2,3,4\}) = 7 + f(\{1,2,3\}) = 8$ and $j = 2$ and $\beta_{8,10} = 2$.



Figure 1

As a final application we give a lower bound for the last nonzero total Betti-number of an unmixed bipartite graph. To describe the result, we introduce the set $B_G \subseteq A_G$ consisting of all elements q such that $|q| - |N(q)| - |\wedge\{r; r \in N(q)\}| = \max_{p \in A_G}\{|p| - |N(p)| - |\wedge\{r; r \in N(p)\}|\}$.

For an R-module M, let $t(M)$ denotes the last nonzero total Betti-number of M. Then we have the following corollary.

Corollary 1.9. *Let G be an unmixed bipartite graph. Then $t(R/I(G)) \geq |B_G|$.*

Proof. Let $p \in B_G$ and $r = \operatorname{pd}(R/I(G))$. Then $r = n + |p| - |N(p)| - |\wedge\{r; r \in N(p)\}|$. Let $b = b(p; N(p))$, from [1, Theorem 2.8] we have $\beta_{r,b}(R/I(G)) = \beta_{|N(p)|,b}(H_{\mathcal{L}_G}) = 1$. Therefore $T(R/I(G)) \geq |B_G|$. □

We do not know of any example of a monomial ideal I of projective dimension r for which there exists a nonzero multigraded Betti-number $\beta_{r,b}$ which is not extremal. If such ideals don't exist, at least among the edge ideals of unmixed bipartite graphs, then we would have equality in Proposition 1.9.

In general the multigraded extremal Betti-numbers of monomial ideal not only appear in the last step of the resolution. However we have

Proposition 1.10. *Let $R = k[x_1, \ldots, x_n]$ for a field k and I a monomial ideal of R such that R/I is a Cohen-Macaulay ring. Then for all multigraded extremal Betti-numbers $\beta_{i,b}(R/I)$ we have $i = \operatorname{pd}(R/I)$.*

Proof. Assume that

$$0 \to F_r \to \cdots \to F_{i+1} \to F_i \to \cdots \to F_0 \to R/I \to 0$$

is a minimal graded free resolution of R/I, where $r = \operatorname{pd}(R/I)$ and φ be the function $F_{i+1} \to F_i$ in the resolution. Let $\beta_{i,b}(R/I)$ be a multigraded extremal Betti-number with $i < \mathrm{pd}(R/I)$ and e be the basis element in F_i with multidegree b. Then for any basis element g of F_{i+1} the coefficient of e in $\varphi(g)$ is zero. Otherwise multideg$(e) <$ multideg(g) and $g \in F_{i+1}$ but then $\beta_{i,b}$ is not extremal, a contradiction. This means that e^* is a cycle in the resolution

$$0 \to (R/I)^* \to F_0^* \to \cdots \to F_i^* \to F_{i+1}^* \to \cdots \to F_r^* \to 0,$$

where $F_i^* = \mathrm{Hom}_R(F_i, R)$. Therefore $\mathrm{Ext}^i_R(R/I, R) \neq 0$. But we know that $\mathrm{Ext}^j_R(R/I, R) = 0$ for any $j < r$, a contradiction. $\square$

References

1. D. Bayer, H. Charalambous, S. Popescu, *Extremal Betti numbers and applications to monomial ideals,* J. Algebra 221 (1999), no. 2, 497–512.
2. H. T. Hà, A. Van Tyul, *Resolutions of squarefree monomial ideals via facet ideals: a survey.* (2006) arXiv preprint math.AC/0604301.
3. H. T. Hà, A. Van Tyul, *Monomial ideals, edge ideals of hypergraphs, and their graded Betti numbers,* J. Algebraic Combin. 27 (2008), no. 2, 215245.
4. J. Herzog, T. Hibi, *Distributive lattices, bipartite graphs and Alexander duality,* J. Alg. Comb. 22 (2005), 289–302.
5. J. Herzog, T. Hibi, H. Ohsugi, *Unmixed bipartite graphs and sublattices of the Boolean lattices,* arXiv preprint math.AC/0806.1088v1 (2008).
6. J. Herzog, T. Hibi, X. Zheng, *Cohen-Macaulay chordal graphs,* (English summary) J. Combin. Theory Ser. A 113 (2006), no. 5, 911–916. 13H10 (05C99)
7. J. Herzog, T. Hibi, X. Zheng, *The monomial ideal of a finite meet-semilattice,* Trans. Amer. Math. Soc. 358 (2006), no. 9, 4119–4134.
8. T. Hibi, *Distributive lattices, affine semigroup rings and algebras with straightening laws,* in "Commutative Algebra and Combinatorics, Advanced Studies in Pure Math." (M. Nagata and H. Matsumura Eds.) Vol. 11, North-Holland, Amesterdam, 1987, 93–109.
9. M. Kummini, *Regularity, Depth and Arithmetic Rank of Bipartite Edge Ideals*, preprint (2008).
10. R. H. Villarreal, *Unmixed bipartite graphs,* To appear in Rev. Colombiana Mat.
11. X. Zheng, *Resolutions of facet ideals,* Comm. Algebra 32 (2004), no. 6, 2301–2324.

Fatemeh Mohammadi, Department of Pure Mathematics, Faculty of Mathematics and Computer Science, Amirkabir University of Technology (Tehran Polytechnic), 424, Hafez Ave., Tehran 15914, Iran.

E-mail address: `f_mohammadi@aut.ac.ir`

Somayeh Moradi, Department of Pure Mathematics, Faculty of Mathematics and Computer Science, Amirkabir University of Technology (Tehran Polytechnic), 424, Hafez Ave., Tehran 15914, Iran.

E-mail address: `s_moradi@aut.ac.ir`